

























How to give the right amount of IBUPROFEN (also known as Motrin, Advil) is different depending on which type of ibuprofen you plan to give.

Dose: Give every 6 hours if needed, for fever or pain. **DO NOT GIVE MORE THAN 4 DOSES IN 24 HOURS.**
Do NOT use with any other medicine containing ibuprofen.

Weight	Age	Infant's Ibuprofen Drops (50 mg / 1.25 mL)	Children's Liquid Ibuprofen (100 mg / 5 mL)	Children's Ibuprofen Chewable Tablets (100 mg)	Adult's Ibuprofen Tablets (200 mg)
0 to 11 pounds (up to 5 kilograms)	0 to 5 months	—	—	—	—
12 to 17 pounds (about 6 to 7 kilograms)	6 to 11 months	1.25 mL 	2.5 mL* 	—	—
18 to 23 pounds (about 8 to 10 kilograms)	12 to 23 months	1.875 mL 	4 mL* 	—	—
24 to 35 pounds (about 11 to 15 kilograms)	2 to 3 years	2.5 mL 	5 mL* → 	1 tablet 	—
36 to 47 pounds (about 16 to 21 kilograms)	4 to 5 years	3.75 mL 	 ← 7.5 mL*	1 ½ tablets 	—
48 to 59 pounds (about 22 to 26 kilograms)	6 to 8 years	5 mL 	10 mL* → 	2 tablets 	1 tablet 
60 to 71 pounds (about 27 to 32 kilograms)	9 to 10 years	—	 ← 12.5 mL*	2 ½ tablets 	1 tablet 
72 to 95 pounds (about 33 to 43 kilograms)	11 years	—	15 mL* → 	3 tablets 	1 ½ tablets 
96 pounds or more (44 kilograms or more)	12 years or older	—	10 mL + 10 mL = 20 mL* 10 mL* →  and 10 mL* → 	4 tablets 	2 tablets 

***Note:** This dosage is for children's liquid ibuprofen products given by dropper. See the previous column for correct dose-by-weight for infant ibuprofen drops, which are more concentrated.



healthychildren.org















Powered by pediatricians. Trusted by parents.
 from the American Academy of Pediatrics

How to give the right amount of DIPHENHYDRAMINE (also known as Benadryl) is different depending on which type of Diphenhydramine you plan to give.

Dose: Give every 6 hours if needed. **DO NOT GIVE MORE THAN 4 DOSES IN 24 HOURS.**

Do **NOT** use with any other medicine with diphenhydramine in it.



Weight →	20 to 24 pounds (about 9 to 10 kilograms)	25 to 37 pounds (about 11 to 16 kilograms)	38 to 49 pounds (about 17 to 22 kilograms)	50 to 99 pounds (about 23 to 45 kilograms)	100 pounds or more (46 kilograms or more)
Children's Liquid Diphenhydramine (12.5 mg / 5 mL)	4 mL 	5 mL 	7.5 mL 	10 mL 	—
Children's Diphenhydramine Chewable Tablets (12.5 mg)	—	1 tablet 	1 ½ tablets 	2 tablets 	4 tablets 
Diphenhydramine Tablets (25 mg)	—	½ tablet 	½ tablet 	1 tablet 	2 tablets 
Diphenhydramine Capsules (25 mg)	—	—	—	1 capsule 	2 capsules 

Do not give to children less than 2 years of age.

Do not give to children 2 to 6 years of age unless your doctor tells you to.


























*TABLE NOTES:

- **Age of child:** Do not give diphenhydramine to children less than 6 years of age unless your child's doctor tells you to. There are other medicines that are like diphenhydramine but will not make your child sleepy (like loratadine, cetirizine, fexofenadine) that can be bought without a prescription and are safer for young children.
- **Measuring the dose for liquid medicines (should be in "mL" or metric units):** It is easier to give the right amount of medicine when using a syringe than when using a kitchen teaspoon or tablespoon. Use the tool that comes with the medicine. If a tool does not come with the medicine, ask your pharmacist for one.
- **How often to give the medicine (frequency):** You can give diphenhydramine every 6 hours as needed.
- **Adult dose:** 50 mg
- **Side effects:** This medicine can make a child sleepy. Some children, however, may get more excited and active instead of getting sleepy. Because this medicine can make people sleepy, it is important to be careful when driving or using heavy machines after taking this medicine. This is especially important for teens who are driving.

How to give the right amount of ACETAMINOPHEN (also known as Tylenol) is different depending on which medicine you plan to give.

Dose: Give every 4 to 6 hours as needed for fever or pain. DO NOT GIVE MORE THAN 4 DOSES IN 24 HOURS.

Do NOT use with any other medicine containing acetaminophen.

Weight	Age	Infant's Acetaminophen (160 mg / 5 mL)	Children's Acetaminophen (160 mg / 5 mL)	Children's Acetaminophen Chewables (160 mg)	Children's Acetaminophen Dissolvable Packets (160 mg / powder pack)	Adult's Acetaminophen Tablets (325 mg)	Adult's Acetaminophen Tablets (500 mg)
6 to 11 pounds (3 to 5 kilograms)	0 to 3 months	1.25 mL 	_____	_____	_____	_____	_____
12 to 17 pounds (about 5 to 7 kilograms)	4 to 11 months	2.5 mL 	_____	_____	_____	_____	_____
18 to 23 pounds (about 8 to 10 kilograms)	12 to 23 months	3.75 mL 	_____	_____	_____	_____	_____
24 to 35 pounds (about 11 to 15 kilograms)	2 to 3 years	_____	5 mL → 	1 tablet 	_____	_____	_____
36 to 47 pounds (about 16 to 21 kilograms)	4 to 5 years	_____	_____ → 7.5 mL 	1 ½ tablets 	_____	_____	_____
48 to 59 pounds (about 22 to 26 kilograms)	6 to 8 years	_____	10 mL → 	2 tablets 	2 packets 	1 tablet 	_____
60 to 71 pounds (about 27 to 32 kilograms)	9 to 10 years	_____	_____ → 12.5 mL 	2 ½ tablets 	2 packets 	1 tablet 	_____
72 to 95 pounds (about 33 to 43 kilograms)	11 years	_____	15 mL → 	3 tablets 	3 packets 	1 ½ tablets 	1 tablet 
96 pounds or more (more than 43 kilograms)	12 years or older	_____	10 mL →  and 10 mL →  10 mL + 10 mL = 20 mL	4 tablets 	_____	2 tablets 	1 tablet 



healthychildren.org

Powered by pediatricians. Trusted by parents.
from the American Academy of Pediatrics